


STRANGE QUARK MATTER 2004

Thermodynamic Properties of Strongly Interacting Matter



A reminder/summary:


QUARK MATTER 2004

FK, J. Phys. G30 (2004) S887 [hep-lat/0403016]

News from Lattice QCD

On


Heavy Quark Potentials and Spectral Functions of Heavy Quark States



Quark Matter 2004 ...

...topics that could not be discussed:


- 1) Update on the QCD phase diagram
 - $T_c(\mu)$ and the chiral critical point: quark mass dependence, lattice artefacts
- 2) QCD equation of state for $\mu > 0$
 - baryon number fluctuations;
 - comparison with resonance gas



Topics from Lattice QCD ...


...to be discussed at SQM2004:


- 1) Update on the QCD phase diagram
 - $T_c(\mu)$ and the chiral critical point: quark mass dependence, lattice artefacts
- 2) QCD equation of state for $\mu > 0$
 - baryon number fluctuations;
 - comparison with resonance gas




Critical behavior in hot and dense matter: QCD phase diagram


crossover vs.
phase transition







Critical behavior in hot and dense matter: QCD phase diagram







Critical behavior in hot and dense matter: QCD phase diagram







Critical behavior in hot and dense matter: QCD phase diagram






Critical behavior in hot and dense matter: QCD phase diagram







Quark mass dependence of the QCD phase diagram






chiral critical point








Quark mass dependence of the QCD phase diagram








Quark mass dependence of the QCD phase diagram






Quark mass dependence of the QCD phase diagram







Quark mass dependence of the QCD phase diagram




- locate critical line at $\mu = 0$





QCD phase diagram for vanishing chemical potential




quark mass dependence


strong cut-off dependence

$m_{PS}^{crit} \simeq 290 \text{ MeV}$, standard staggered
 $m_{PS}^{crit} \simeq 70 \text{ MeV}$, improved staggered
(Bielefeld, hep-lat/0309077)

- continuous transition for a wide range of quark masses



QCD phase diagram for vanishing chemical potential



- continuous transition for a wide range of quark masses

$$n_f = 3:$$


- critical point in Ising universality class

expansion around $m_s = m_{u,d}$:


$$m_s^{crit} = 3m_{u,d}^{crit,3f} - 2m_{u,d}$$




$$(m_K^{crit})^2 = \frac{3}{2} (m_\pi^{crit,3f})^2 - \frac{1}{2} m_\pi^2$$




QCD phase diagram for vanishing chemical potential




- continuous transition for a wide range of quark masses
- curvature of critical surface at $\mu = 0$ for $n_f = 3$ not crucial







QCD phase diagram for vanishing chemical potential




- continuous transition for a wide range of quark masses
- curvature of critical surface at $\mu = 0$ for $n_f = 3$ not crucial






The chiral condensate: 2- vs. (2+1)- vs. 3-flavor QCD





- compare simulations in 3- and (2+1)-flavor QCD

C. Bernard et al. (MILC Col.), hep-lat/0405029



The chiral condensate: 2- vs. (2+1)- vs. 3-flavor QCD



- compare simulations in 3- and (2+1)-flavor QCD


- $\langle \bar{\psi} \psi \rangle$ for 3- and (2+1)-flavor QCD almost identical;

- sensitive only to light quark mass


C. Bernard et al. (MILC Col.), hep-lat/0405029



chiral properties of 2-flavor QCD dominate (pseudo)-critical behavior



The chiral condensate: 2- vs. (2+1)- vs. 3-flavor QCD



$n_f = 2$:

- O(4) universality class expected, **but:**
could not yet be verified

FK, E. Laermann, PRD50 (1994) 6954


doubts:

evidence for 1st order transition??

A. DiGiacomo et al., hep-lat/0408008



chiral properties of 2-flavor QCD dominate (pseudo)-critical behavior




Extending the phase diagram to non-vanishing chemical potential

non-zero baryon number density: $\mu > 0$

$$\begin{aligned} Z(\mathbf{V}, \mathbf{T}, \boldsymbol{\mu}) &= \int \mathcal{D}\mathcal{A} \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-S_E(\mathbf{V}, \mathbf{T}, \boldsymbol{\mu})} \\ &= \int \mathcal{D}\mathcal{A} \mathcal{D} \det M(\boldsymbol{\mu}) e^{-S_E(\mathbf{V}, \mathbf{T})} \end{aligned}$$

↑complex fermion determinant;
long standing problem

⇒ three (partial) solutions for large T , small μ




Extending the phase diagram to non-vanishing chemical potential

non-zero baryon number density: $\mu > 0$

$$\begin{aligned} Z(\mathbf{V}, \mathbf{T}, \boldsymbol{\mu}) &= \int \mathcal{D}\mathcal{A} \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-S_E(\mathbf{V}, \mathbf{T}, \boldsymbol{\mu})} \\ &= \int \mathcal{D}\mathcal{A} \mathcal{D} \det M(\boldsymbol{\mu}) e^{-S_E(\mathbf{V}, \mathbf{T})} \end{aligned}$$

\uparrow complex fermion determinant;
long standing problem

- ⇒ three (partial) solutions for large T , small μ
- exact evaluation of $\det M$: works well on small lattices; requires reweighting
[Z. Fodor, S.D. Katz, JHEP 0203 \(2002\) 014](#)
- Taylor expansion around $\mu = 0$ works well for small μ ; has truncation errors
[C. R. Allton et al. \(Bielefeld-Swansea\), Phys. Rev. D66 \(2002\) 074507](#)
- imaginary chemical potential: works well for small μ ; requires analytic continuation
[Ph. deForcrand, O. Philipsen, Nucl. Phys. B642 \(2002\) 290](#)



Extending the phase diagram to non-vanishing chemical potential

non-zero baryon number density: $\mu > 0$


$$\begin{aligned} Z(\mathbf{V}, \mathbf{T}, \boldsymbol{\mu}) &= \int \mathcal{D}\mathcal{A} \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-S_E(\mathbf{V}, \mathbf{T}, \boldsymbol{\mu})} \\ &= \int \mathcal{D}\mathcal{A} \mathcal{D} \det M(\boldsymbol{\mu}) e^{-S_E(\mathbf{V}, \mathbf{T})} \end{aligned}$$

↑complex fermion determinant;
long standing problem

⇒ three (partial) solutions for large T , small μ

$$T_c(\mu) = ? \quad \boxed{\frac{dT_c}{d(\mu)^2} = -\frac{1}{N_\tau^2 T_c(0)} \frac{\partial g_c^{-2}(\mu a)}{\partial (\mu a)^2} \left(a \frac{\partial g^{-2}(a)}{\partial a} \right)}$$

- determine μ -dependence of Lee-Yang zeroes of partition function
- determine shift of peak positions of susceptibilities and/or radius of convergence of Taylor series
- locate phase transition (crossover) for imaginary μ



Extending the phase diagram to non-vanishing chemical potential

non-zero baryon number density: $\mu > 0$

$$\begin{aligned} Z(\mathbf{V}, \mathbf{T}, \boldsymbol{\mu}) &= \int \mathcal{D}\mathcal{A} \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-S_E(\mathbf{V}, \mathbf{T}, \boldsymbol{\mu})} \\ &= \int \mathcal{D}\mathcal{A} \mathcal{D} \det M(\boldsymbol{\mu}) e^{-S_E(\mathbf{V}, \mathbf{T})} \end{aligned}$$


pert. or non-pert. β -function

$T_c(\mu) = ?$

$$\boxed{\frac{dT_c}{d(\mu)^2} = -\frac{1}{N_\tau^2 T_c(0)} \frac{\partial g_c^{-2}(\mu a)}{\partial (\mu a)^2} \left(a \frac{\partial g^{-2}(a)}{\partial a} \right)}$$




- determine μ -dependence of Lee-Yang zeroes of partition function
- determine shift of peak positions of susceptibilities and/or radius of convergence of Taylor series
- locate phase transition (crossover) for imaginary μ




Reweighting of susceptibilities for small μ

Polyakov loop susceptibility




chiral susceptibility




$$T_c(\mu_B)/T_c(0) = 1 - 0.0078(38) (\mu_B/T_c(0))^2 \quad (\text{non-pert.})$$

C. R. Allton et al. (Bielefeld-Swansea), Phys. Rev. D66 (2002) 074507



Extending the phase diagram to non-vanishing chemical potential

analysis of volume dependence of Lee-Yang zeroes for $\mu > 0$



Fodor & Katz,
JHEP 0203 (2002) 014


$V = 4^3, 6^3, 8^3$

Fodor & Katz,
JHEP 0404 (2004) 050

$V = 6^3, 8^3, 10^3, 12^3$


$$\mu^{crit} = 360(40) \text{ MeV}$$

$$T_c(\mu_B)/T_c(0) = 1 - 0.0032(1) (\mu_B/T_c(0))^2 \quad (\text{pert.})$$



Extending the phase diagram to non-vanishing chemical potential

first (exploratory) results on the quark mass dependence of the transition line:




m_q -dependence

(3-flavor QCD, pert. β -function, Taylor expansion)

$$\frac{T_c(\mu)}{T_c(0)} : \begin{aligned} &1 - 0.025(6)(\mu_q/T)^2, \text{ } ma = 0.1 \\ &1 - 0.114(46)(\mu_q/T)^2, \text{ } ma = 0.005 \end{aligned}$$


Bielefeld-Swansea

(hep-lat/0309116, Lattice 2003)



Extending the phase diagram to non-vanishing chemical potential

first (exploratory) results on the quark mass dependence of the transition line:



m_q -dependence not confirmed in simulations with imaginary μ

Ph. de Forcrand, O. Philipsen, NP B673 (2003) 170

m_q -dependence


(3-flavor QCD, pert. β -function, Taylor expansion)

$$\frac{T_c(\mu)}{T_c(0)} : \begin{aligned} & 1 - 0.025(6)(\mu_q/T)^2, \quad ma = 0.1 \\ & 1 - 0.114(46)(\mu_q/T)^2, \quad ma = 0.005 \end{aligned}$$

Bielefeld-Swansea

(hep-lat/0309116, Lattice 2003)


a systematic analysis of
cut-off effects, scaling violations
AND volume + truncation effects
still needs to be done




Extending the phase diagram to non-vanishing chemical potential

non-zero baryon number density: $\mu > 0$

$$\begin{aligned} Z(\mathbf{V}, \mathbf{T}, \boldsymbol{\mu}) &= \int \mathcal{D}\mathcal{A} \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-S_E(\mathbf{V}, \mathbf{T}, \boldsymbol{\mu})} \\ &= \int \mathcal{D}\mathcal{A} \mathcal{D} \det M(\boldsymbol{\mu}) e^{-S_E(\mathbf{V}, \mathbf{T})} \end{aligned}$$




$$\frac{T_c(\mu)}{T_c(0)} : \begin{aligned} &1 - 0.0056(4)(\mu_B/T)^2 \\ &\text{deForcrand, Philipsen (imag. } \mu, \text{ pert)} \\ &1 - 0.0078(38)(\mu_B/T)^2 \\ &\text{Bielefeld-Swansea} \\ &(\mathcal{O}(\mu^2) \text{ reweighting, non-pert)} \\ &1 - 0.0032(1)(\mu_B/T)^2 \\ &\text{Fodor,Katz(Lee-Yang zeroes, pert)} \end{aligned}$$




Extending the phase diagram to non-vanishing chemical potential

non-zero baryon number density: $\mu > 0$

$$\begin{aligned} Z(\mathbf{V}, \mathbf{T}, \boldsymbol{\mu}) &= \int \mathcal{D}\mathcal{A} \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-S_E(\mathbf{V}, \mathbf{T}, \boldsymbol{\mu})} \\ &= \int \mathcal{D}\mathcal{A} \mathcal{D} \det M(\boldsymbol{\mu}) e^{-S_E(\mathbf{V}, \mathbf{T})} \end{aligned}$$




current studies of $T_c(\mu)$ are exploratory!
uncertainties in scale-determination and
systematics of quark mass dependenceee



Extending the phase diagram to non-vanishing chemical potential

non-zero baryon number density: $\mu > 0$

$$\begin{aligned} Z(\mathbf{V}, \mathbf{T}, \boldsymbol{\mu}) &= \int \mathcal{D}\mathbf{A} \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-S_E(\mathbf{V}, \mathbf{T}, \boldsymbol{\mu})} \\ &= \int \mathcal{D}\mathbf{A} \det M(\boldsymbol{\mu}) e^{-S_E(\mathbf{V}, \mathbf{T})} \end{aligned}$$



current studies of $T_c(\mu)$ are exploratory!

uncertainties in scale-determination and
systematics of quark mass dependenceee


GSI future

RHIC, LHC

$T_c(\mu) \equiv T_{\text{freeze}}$?

P. Braun-Munzinger, J. Stachel,
C. Wetterich, hep-nucl/0311005

Will be answered by LGT calculations




Intermezzo: Next generation computers for lattice gauge theory



today:

computing resources not sufficient
for reliable thermodynamics calculations
with physical quark mass parameters,
 $\mu \neq 0$ and large enough statistics
on large enough lattices

120 GFlops APEmille in Bielefeld




Intermezzo: Next generation computers for lattice gauge theory

QCDOC and apeNEXT

2004/05:



QCD thermodynamics on the next generation of special purpose
dedicated QCD computers


installations with (10-20) TFlops peak speed are planned
in the USA and Europe



apeNEXT: Next generation of APE computers


Assembling apeNEXT...






apeNEXT: Next generation of APE computers


Assembling apeNEXT...






apeNEXT: Next generation of APE computers


Assembling apeNEXT...



- first chips Dec. 2003
- two 0.8 TFlops prototypes ~ autumn 2004
- first 3 TFlops installations in 2005



BackPlane



QCDOC: Next generation of Columbia-RIKEN computer

Columbia – RIKEN – UKQCD Collaboration




512 – node machine : (360 – 450) GFlops

- currently debugging three 0.9-Teraflops machines (09/2004):
1024-node systems, 0.5 Tbyte memory; 6 Gbit/sec Ethernet I/O bandwidth


QCDOC computing center at BNL :

- 10 TFlops machine for RBRC: ~ autumn 2004
- 10 TFlops machine for american LGT community: ~ early 2005
- ... larger installations possible and needed!



Bulk thermodynamics with non-vanishing chemical potential

$$\begin{aligned} Z(\mathbf{V}, \mathbf{T}, \boldsymbol{\mu}) &= \int \mathcal{D}\mathcal{A} \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-S_E(\mathbf{V}, \mathbf{T}, \boldsymbol{\mu})} \\ &= \int \mathcal{D}\mathcal{A} [\det M(\boldsymbol{\mu})]^f e^{-S_G(\mathbf{V}, \mathbf{T})} \\ &\quad \uparrow \text{complex fermion determinant}; \end{aligned}$$




Bulk thermodynamics with non-vanishing chemical potential

$$\begin{aligned} Z(\mathbf{V}, \mathbf{T}, \boldsymbol{\mu}) &= \int \mathcal{D}\mathcal{A} \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-S_E(\mathbf{V}, \mathbf{T}, \boldsymbol{\mu})} \\ &= \int \mathcal{D}\mathcal{A} [\det M(\boldsymbol{\mu})]^f e^{-S_G(\mathbf{V}, \mathbf{T})} \end{aligned}$$


↑complex fermion determinant;
↓Taylor expansion;

$$\begin{aligned} \frac{p}{T^4} &= \frac{1}{VT^3} \ln Z(\mathbf{V}, \mathbf{T}, \boldsymbol{\mu}) \\ &\equiv \sum_{n=0}^{\infty} c_n(T) \left(\frac{\mu}{T} \right)^n \\ &= c_0 + c_2 \left(\frac{\mu}{T} \right)^2 + c_4 \left(\frac{\mu}{T} \right)^4 + \mathcal{O}((\mu/T)^6) \end{aligned}$$

$$\boldsymbol{\mu} = \mathbf{0} \quad \Rightarrow \quad \frac{p}{T^4} \equiv c_0(T)$$



$\mu \equiv 0$: T_c and equation of state



$$T_c = (173 \pm 8 \pm \text{sys}) \text{ MeV}$$

FK, E. Laermann, A. Peikert,
Nucl. Phys. B605 (2001) 579


$$T_c = 167(13) [177(11)] \text{ MeV}$$


C. Bernard et al. (MILC Col.), hep-lat/0405029

$$\epsilon_c \simeq (6 \pm 2) T_c^4$$


$$\simeq (0.3 - 1.3) \text{ GeV/fm}^3$$

energy density for 0, 2 and 3-flavor QCD





$\mu \equiv 0$: T_c and equation of state



$$\epsilon_c \simeq (6 \pm 2) T_c^4$$
$$\simeq (0.3 - 1.3) \text{ GeV/fm}^3$$

Can we be satisfied
with these results?


$$T_c = (173 \pm 8 \pm \text{sys}) \text{ MeV}$$


FK, E. Laermann, A. Peikert,
Nucl. Phys. B605 (2001) 579

$$T_c = 167(13) [177(11)] \text{ MeV}$$


C. Bernard et al. (MILC Col.), hep-lat/0405029

Energy density for 0, 2 and 3-flavor QCD





$\mu \equiv 0$: T_c and equation of state




T_c


- $m_{PS} \gtrsim 300 \text{ MeV}$ (chiral limit??)
- $a \simeq 0.2 \text{ fm}$ (continuum limit??)
- improved staggered fermions,
⇒ flavor symmetry breaking
(need even better fermion actions)

ϵ_c

- $m_{PS} \simeq 770 \text{ MeV}$ (!!?)
- $V \simeq (4 \text{ fm})^3$ (thermodynamic limit)

energy density for 0, 2 and 3-flavor QCD





Pressure etc. up to $\mathcal{O}((\mu_q/T)^6)$

- Taylor expansion up to $\mathcal{O}((\mu_q/T)^6)$

$$\frac{p}{T^4} = \frac{1}{VT^3} \ln \mathcal{Z} = \sum_{n=0}^{\infty} c_n(T) \left(\frac{\mu_q}{T} \right)^n \simeq c_0 + c_2 \left(\frac{\mu_q}{T} \right)^2 + c_4 \left(\frac{\mu_q}{T} \right)^4 + c_6 \left(\frac{\mu_q}{T} \right)^6$$


quark number density $\frac{n_q}{T^3} = 2c_2 \frac{\mu_q}{T} + 4c_4 \left(\frac{\mu_q}{T} \right)^3 + 6c_6 \left(\frac{\mu_q}{T} \right)^5$

quark number susceptibility $\frac{\chi_q}{T^2} = 2c_2 + 12c_4 \left(\frac{\mu_q}{T} \right)^2 + 30c_6 \left(\frac{\mu_q}{T} \right)^4$

an **estimator** for the radius of convergence

$$\left(\frac{\mu_q}{T} \right)_{crit} = \lim_{n \rightarrow \infty} \left| \frac{c_{n-2}}{c_n} \right|^{1/2}$$




$c_n > 0$ for all n ;
singularity for real μ




Pressure etc. up to $\mathcal{O}((\mu_q/T)^6)$

- Taylor expansion up to $\mathcal{O}((\mu_q/T)^6)$

$$\frac{p}{T^4} = \frac{1}{VT^3} \ln \mathcal{Z} = \sum_{n=0}^{\infty} c_n(T) \left(\frac{\mu_q}{T} \right)^n \simeq c_0 + c_2 \left(\frac{\mu_q}{T} \right)^2 + c_4 \left(\frac{\mu_q}{T} \right)^4 + c_6 \left(\frac{\mu_q}{T} \right)^6$$







Pressure etc. up to $\mathcal{O}((\mu_q/T)^6)$

- Taylor expansion up to $\mathcal{O}((\mu_q/T)^6)$

$$\frac{p}{T^4} = \frac{1}{VT^3} \ln \mathcal{Z} = \sum_{n=0}^{\infty} c_n(T) \left(\frac{\mu_q}{T} \right)^n \simeq c_0 + c_2 \left(\frac{\mu_q}{T} \right)^2 + c_4 \left(\frac{\mu_q}{T} \right)^4 + c_6 \left(\frac{\mu_q}{T} \right)^6$$




$c_n > 0$ for all n and $T \lesssim 0.95 T_c \Leftrightarrow$ singularity for real μ (positive μ^2)




Pressure etc. up to $\mathcal{O}((\mu_q/T)^6)$

- Taylor expansion up to $\mathcal{O}((\mu_q/T)^6)$

$$\frac{p}{T^4} = \frac{1}{VT^3} \ln \mathcal{Z} = \sum_{n=0}^{\infty} c_n(T) \left(\frac{\mu_q}{T} \right)^n \simeq c_0 + c_2 \left(\frac{\mu_q}{T} \right)^2 + c_4 \left(\frac{\mu_q}{T} \right)^4 + c_6 \left(\frac{\mu_q}{T} \right)^6$$



irregular sign of c_n for $T \gtrsim T_c$ \Leftrightarrow singularity in complex plane




Radius of convergence: lattice estimates vs. resonance gas




Taylor expansion \Rightarrow estimates for radius of convergence


$$\rho_n = \sqrt{\left| \frac{c_{2n-2}}{c_{2n}} \right|}$$



$T < T_0$: $0.8 \lesssim \rho_n \lesssim 1.5$ for all n



$\Rightarrow 400 \text{ MeV} \lesssim \mu_B^{crit} \lesssim 800 \text{ MeV}$





Radius of convergence: lattice estimates vs. resonance gas




Taylor expansion \Rightarrow estimates for radius of convergence

$$\rho_n = \sqrt{\left| \frac{c_{2n-2}}{c_{2n}} \right|}$$



$T < T_0: 0.8 \lesssim \rho_n \lesssim 1.5$ for all $n \Rightarrow 400 \text{ MeV} \lesssim \mu_B^{crit} \lesssim 800 \text{ MeV}$


HOWEVER: see discussion of resonance gas!!!



The pressure for $\mu_q/T > 0$

C.R. Allton et al. (Bielefeld-Swansea), PRD68 (2003) 014507


$\mu_q = 0$, $16^3 \times 4$ lattice
improved staggered fermions;
 $n_f = 2$, $m_\pi \simeq 770 \text{ MeV}$




high-T, ideal gas limit

$$\frac{p}{T^4} \Big|_\infty = n_f \left(\frac{7\pi^2}{60} + \frac{1}{2} \left(\frac{\mu_q}{T} \right)^2 + \frac{1}{4\pi^2} \left(\frac{\mu_q}{T} \right)^4 \right)$$

contribution from $\mu_q/T > 0$
Taylor expansion, $\mathcal{O}((\mu/T)^4)$






The pressure for $\mu_q/T > 0$


C.R. Allton et al. (Bielefeld-Swansea), PRD68 (2003) 014507


$\mu_q = 0$, $16^3 \times 4$ lattice
improved staggered fermions;
 $n_f = 2$, $m_\pi \simeq 770$ MeV



pattern for $\mu_q = 0$ and $\mu_q > 0$ similar;
quite large contribution in hadronic phase;
 $\mathcal{O}((\mu/T)^6)$ correction small for $\mu_q/T \lesssim 1$

contribution from $\mu_q/T > 0$
NEW: Taylor expansion, $\mathcal{O}((\mu/T)^6)$






Fluctuations of the quark number density ($\mu_q > 0$)

quark number density fluctuations:

up to $\mathcal{O}((\mu_q/T)^2)$



$$\frac{\chi_q}{T^2} = \left(\frac{\partial^2}{\partial(\mu_q/T)^2} \frac{p}{T^4} \right)_{T \text{ fixed}}$$

high-T, massless limit: polynomial in (μ_q/T)


$$\frac{\chi_{q,SB}}{T^2} = n_f + \frac{3n_f}{\pi^2} \left(\frac{\mu_q}{T} \right)^2$$

large density fluctuations closer to the chiral critical point

$$\frac{\chi_q}{T^2} = \frac{1}{VT^3} \left(\langle N_q^2 \rangle - \langle N_q \rangle^2 \right)$$


$$\left(\frac{\partial p}{\partial n_q} \right)_T = \frac{n_q}{\chi_q}$$

$\Rightarrow \chi_q$ will diverge on chiral critical point



Fluctuations of the quark number density ($\mu_q > 0$)

quark number density fluctuations:
up to $\mathcal{O}((\mu_q/T)^4)$



$$\frac{\chi_q}{T^2} = \left(\frac{\partial^2}{\partial(\mu_q/T)^2} \frac{p}{T^4} \right)_{T \text{ fixed}}$$

high-T, massless limit: polynomial in (μ_q/T)


$$\frac{\chi_{q,SB}}{T^2} = n_f + \frac{3n_f}{\pi^2} \left(\frac{\mu_q}{T} \right)^2$$

large density fluctuations closer to the chiral critical point

$$\frac{\chi_q}{T^2} = \frac{1}{VT^3} \left(\langle N_q^2 \rangle - \langle N_q \rangle^2 \right)$$

$$\left(\frac{\partial p}{\partial n_q} \right)_T = \frac{n_q}{\chi_q}$$



$\Rightarrow \chi_q$ will diverge on chiral critical point




Fluctuations of the quark number density ($\mu_q > 0$)

quark number density fluctuations:


$$\frac{\chi_q}{T^2} = \left(\frac{\partial^2}{\partial(\mu_q/T)^2} \frac{p}{T^4} \right)_{T \text{ fixed}}$$





Fluctuations of the isospin and charge densities ($\mu_q > 0$, $\mu_I = 0$)

isospin density fluctuations:
up to $\mathcal{O}((\mu_q/T)^2)$




$$\frac{\chi_I}{T^2} = \left(\frac{\partial^2}{\partial(\mu_I/T)^2} \frac{p}{T^4} \right)_{T, \mu_q \text{ fixed}}$$

$\mu_I = 0$:

high-T, massless limit: polynomial in (μ_q/T)

$$\frac{\chi_{I,SB}}{T^2} \equiv \frac{\chi_{q,SB}}{T^2} = n_f + \frac{3n_f}{\pi^2} \left(\frac{\mu_q}{T} \right)^2$$


$$\frac{\chi_{I,SB}}{T^2} = \frac{1}{VT^3} \langle (N_u - N_d)^2 \rangle$$



Fluctuations of the isospin and charge densities ($\mu_q > 0$, $\mu_I = 0$)

isospin density fluctuations:

up to $\mathcal{O}((\mu_q/T)^4)$




$$\frac{\chi_I}{T^2} = \left(\frac{\partial^2}{\partial(\mu_I/T)^2} \frac{p}{T^4} \right)_{T, \mu_q \text{ fixed}}$$

$\mu_I = 0$:

high-T, massless limit: polynomial in (μ_q/T)

$$\frac{\chi_{I,SB}}{T^2} \equiv \frac{\chi_{q,SB}}{T^2} = n_f + \frac{3n_f}{\pi^2} \left(\frac{\mu_q}{T} \right)^2$$


$$\frac{\chi_{I,SB}}{T^2} = \frac{1}{VT^3} \langle (N_u - N_d)^2 \rangle$$



Fluctuations of the isospin and charge densities ($\mu_q > 0$, $\mu_I = 0$)


baryon number density and isospin density fluctuations

charge fluctuations



$$\chi_Q = \frac{1}{36} \chi_q + \frac{1}{4} \chi_I$$


$\sim \chi_q$ close to $T_c(\mu_c)$



Fluctuations of the isospin and charge densities ($\mu_q > 0, \mu_I = 0$)

baryon number density and isospin density fluctuations

charge fluctuations




$$\chi_Q = \frac{1}{36} \chi_q + \frac{1}{4} \chi_I$$

$$\sim \chi_q \quad \text{close to } T_c(\mu_c)$$

for comparison: proton number fluctuations;
 $m_p/T_0 \simeq 5.5$, ideal gas
 \Rightarrow for $T \equiv T_0$, $\mu_q/T = 1$:

$$(\chi_q/T^2)^{\text{proton}} \simeq 1.2 \ll \chi_q/T^2$$



Critical behaviour in dense matter

Hagedorn's Resonance Gas

R. Hagedorn, Nuovo Cimento 35 (1965) 395

strong interactions \Rightarrow exponentially rising spectrum of resonances

Hagedorn spectrum : $\rho(m_H) \sim c m_H^a e^{m_H/T_H}$


$$Z(T, \mu_B) = \int dm_H \rho(m_H) e^{-m_H/T}$$



\Rightarrow critical behaviour: $T_c \equiv T_H$
(end of hadronic physics)

$$\rho(m_H) \equiv \rho_{\text{meson}}(m_H, T) + \rho_{\text{baryon}}(m_H, T, \mu_B)$$

- $\int \Rightarrow \sum$ experimentally known resonances
- $\rho(m_H) \Rightarrow \rho^{\text{lat}}(m_H(m_\pi))$



Critical behaviour in dense matter


Hagedorn's Resonance Gas

R. Hagedorn, Nuovo Cimento 35 (1965) 395

strong interactions \Rightarrow exponentially rising spectrum of resonances

Hagedorn spectrum : $\rho(m_H) \sim c m_H^a e^{m_H/T_H}$

$$Z(T, \mu_B) = \int dm_H \rho(m_H) e^{-m_H/T}$$



\Rightarrow critical behaviour: $T_c \equiv T_H$
(end of hadronic physics)

resonance gas (adjusted mass spectrum):
 ~ 1000 resonance d.o.f.


lattice calculation:

(2+1)-flavor QCD, $m_q/T = 0.4$, $\mu_B = 0$

continuous transition at T_c

FK, K. Redlich, A. Tawfik, hep-ph/0303108

F. Karsch, Strange Quark Matter 2004 – p.25/35



Baryonic resonance gas ⇒ Boltzmann approximation

heavy resonances, $T \ll m_H \Rightarrow$ Boltzmann statistics

$$\mu_B \equiv 3\mu_q$$

thermodynamics: $p(T, \mu_B) = \frac{T}{V} \ln Z(T, \mu_B, V) = \sum_m p_m(T, \mu_B)$

$$\ln Z(T, \mu_B, V) = \sum_{i \in \text{mesons}} \ln Z_{m_i}^B(T, V) + \sum_{i \in \text{baryons}} \ln Z_{m_i}^F(T, \mu_B, V)$$

contribution of baryons with mass m

$$\frac{p_m}{T^4} = \frac{d}{\pi^2} \left(\frac{m}{T}\right)^2 \sum_{\ell=1}^{\infty} (-1)^{\ell+1} \ell^{-2} K_2(\ell m/T) \cosh(\ell \mu_B/T)$$




$$K_2(x) \simeq \sqrt{\pi/2x} \exp(-x), \quad x \gg 1$$

- only $\ell = 1$ contributes for $(m_N - \mu_B) \gtrsim T$

⇒ Boltzmann approximation:

$$\frac{p_m}{T^4} = \frac{d}{\pi^2} \left(\frac{m}{T}\right)^2 K_2(m/T) \cosh(\mu_B/T)$$




Boltzmann approximation with and without Taylor expansion

- Boltzmann approximation \Rightarrow factorization of T and (μ/T) -dependence

		full result	Taylor expansion
pressure	$\frac{\Delta p}{T^4}$	$F(T) [\cosh \frac{\mu_B}{T} - 1]$	$F(T) \left(\tilde{c}_2 \left(\frac{\mu_q}{T} \right)^2 + \tilde{c}_4 \left(\frac{\mu_q}{T} \right)^4 \right)$
number density	$\frac{n_q}{T^3}$	$3F(T) \sinh \frac{\mu_B}{T}$	$F(T) \left(2\tilde{c}_2 \left(\frac{\mu_q}{T} \right) + 4\tilde{c}_4 \left(\frac{\mu_q}{T} \right)^3 \right)$
quark number	$\frac{\chi_q}{T^2}$	$9F(T) \cosh \frac{\mu_B}{T}$	$F(T) \left(2\tilde{c}_2 + 12\tilde{c}_4 \left(\frac{\mu_q}{T} \right)^2 \right)$
susceptibility			

- $c_n = F(T) \tilde{c}_n$, with $\tilde{c}_2 = 9/2$, $\tilde{c}_4 = 27/8$ and $F(T) = \sum_i \frac{d_i}{\pi^2} \left(\frac{m_i}{T} \right)^2 K_2(m_i/T)$
- Taylor expansion only depends on $c_2(T)$ and spectrum independent ratios: c_n/c_2






Resonance gas: spectrum independent consequences

FK, K. Redlich, A. Tawfik, PLB 571 (2003) 67

- Boltzmann approximation \Rightarrow factorization \Rightarrow temperature independent ratios; spectrum independent results

$$\frac{\Delta p}{T^2 \chi_q} = \frac{1}{9} \left(1 - \cosh^{-1}(3\mu_q/T) \right) \sim \frac{\left(\frac{\mu_q}{T}\right)^2 + \frac{c_4}{c_2} \left(\frac{\mu_q}{T}\right)^4 + \frac{c_6}{c_2} \left(\frac{\mu_q}{T}\right)^6}{2 + 12\frac{c_4}{c_2} \left(\frac{\mu_q}{T}\right)^2 + 30\frac{c_6}{c_2} \left(\frac{\mu_q}{T}\right)^4}$$








Resonance gas: spectrum independent consequences

FK, K. Redlich, A. Tawfik, PLB 571 (2003) 67

- Boltzmann approximation \Rightarrow factorization \Rightarrow temperature independent ratios; spectrum independent results

$$\frac{\Delta p}{T^2 \chi_q} = \frac{1}{9} \left(1 - \cosh^{-1}(3\mu_q/T) \right) \sim \frac{\left(\frac{\mu_q}{T}\right)^2 + \frac{c_4}{c_2} \left(\frac{\mu_q}{T}\right)^4 + \frac{c_6}{c_2} \left(\frac{\mu_q}{T}\right)^6}{2 + 12\frac{c_4}{c_2} \left(\frac{\mu_q}{T}\right)^2 + 30\frac{c_6}{c_2} \left(\frac{\mu_q}{T}\right)^4}$$






Resonance gas: spectrum dependent consequences




"fit" with modified spectrum
 \Rightarrow tests factorization

$$m_H(m_\pi) = m_H(0) + A \left(\frac{m_\pi}{m_H(0)} \right)^2$$


$$\frac{\chi_q}{T^2} = 9F(T) \cosh(3\mu_q/T) \sim c_2(T) \left(2 + 12 \frac{c_4}{c_2} \left(\frac{\mu_q}{T} \right)^2 + \mathcal{O} \left(\left(\frac{\mu_q}{T} \right)^4 \right) \right)$$



$$A = 0.9, 1.0, 1.1, 1.2$$



$$A = 1.0$$



Critical behaviour and the resonance gas

- Resonance gas \Rightarrow no critical behaviour, analytic, infinite radius of convergence
 - finite density QCD \Rightarrow expect 1st order phase transition for

$$T < T_c(\mu_c), \quad \frac{\mu_c}{T_c} \sim \mathcal{O}(1);$$

signaled by density fluctuations, i.e.


$\chi_q \rightarrow \infty$ for $\mu \rightarrow \mu_c$ at $T = T_c(\mu_c)$,

resonance gas approximation **should break down** close to the transition line;

1) $\frac{n_q}{T\chi_q} \rightarrow 0$ and $\frac{\Delta p}{T^2\chi_q} \rightarrow 0$ at the chiral critical point NO

2) $|c_{2n-2}/c_{2n}|$ should stay $\mathcal{O}(1)$ YES


SO FAR: no clear evidence for break-down of the resonance gas approximation below T_c



Strange Quark Matter 2004 ...

...topics that could not be discussed:

- 1) Heavy quark spectral functions
 - quarkonium suppression at high temperature
- 2) Heavy quark free energies and the heavy quark potential
 - renormalization of free energies;
 - running coupling at finite temperature






Heavy quark spectral functions comparison of different approaches

M. Asakawa, T. Hatsuda, hep-lat/0308034

S. Datta et al., hep-lat/0312037

J/ψ spectral function







Heavy quark spectral functions comparison of different approaches

M. Asakawa, T. Hatsuda, hep-lat/0308034


J/ψ spectral function




J/ψ dissociates for $1.6T_c \lesssim T \lesssim 1.9T_c$
rather abrupt disappearance of *J/ψ*




S. Datta et al., hep-lat/0312037



J/ψ gradually disappears for $T \gtrsim 1.5T_c$
J/ψ strength reduced by 25% at $T = 2.25T_c$







Heavy quark spectral functions comparison of different approaches

M. Asakawa, T. Hatsuda, hep-lat/0308034


J/ψ spectral function




J/ψ dissociates for $1.6T_c \lesssim T \lesssim 1.9T_c$
rather abrupt disappearance of *J/ψ*




S. Datta et al., hep-lat/0312037



J/ψ gradually disappears for $T \gtrsim 1.5T_c$
J/ψ strength reduced by 25% at $T = 2.25T_c$



radial excitations (χ_c) disappear
already close to T_c



Color averaged heavy quark free energies


O. Kaczmarek, FK, P. Petreczky, F.Zantow, PLB 543(2002)41


confined phase ($T < T_c$) :

deconfined phase ($T > T_c$) :

matching (\equiv renormalization) at short distances to $T = 0$ potential for all T

$$F = E - TS$$





Running coupling from singlet free energy


O. Kaczmarek, FK, P. Petreczky, F. Zantow, hep-lat/0406036

- singlet free energy defines a running coupling:

$$\alpha_{qq} = \frac{3r^2}{4} \frac{dF_1(r, T)}{dr}$$


large distance: constant Coulomb term (string model)

short distance: running coupling
 $\alpha(r)$ from ($T = 0$), 3-loop
(S. Necco, R. Sommer,
Nucl. Phys. B622 (2002) 328)



- short distance physics \Leftrightarrow vacuum physics

T-dependence starts in non-perturbative regime for $T \lesssim 3 T_c$



Conclusions

- QCD thermodynamics for small μ and large T can be analyzed reliably
 - results on bulk thermodynamics in the hadronic phase so far agree quite well with resonance gas model calculations
- rapid rise of quark number fluctuations with increasing density (μ/T) is "normal" in the hadronic phase
 - even larger fluctuations should signal presence of the chiral critical point;
 - higher orders in the Taylor expansion needed;
 - large density fluctuations call for large lattices
- location of the chiral critical point and curvature of $T_c(\mu)$ is uncertain to at least within a factor of 2.
 - would like to see direct evidence for the first order transition

direct simulations at $\mu > 0 ???$